

Accelerated Article Preview

Low neutralization of SARS-CoV-2 Omicron BA.2.75.2, BQ.1.1, and XBB.1 by parental mRNA vaccine or a BA.5-bivalent booster

Received: 31 October 2022

Accepted: 30 November 2022

Accelerated Article Preview

Published online: 06 December 2022

Cite this article as: Kurhade, C. . et al. Low neutralization of SARS-CoV-2 Omicron BA.2.75.2, BQ.1.1, and XBB.1 by parental mRNA vaccine or a BA.5-bivalent booster. *Nature Medicine* <https://doi.org/10.1038/s41591-022-02162-x> (2022).

Chaitanya Kurhade, Jing Zou, Hongjie Xia, Mingru Liu, Hope C. Chang, Ping Ren, Xuping Xie & Pei-Yong Shi

This is a PDF file of a peer-reviewed paper that has been accepted for publication.

Although unedited, the content has been subjected to preliminary formatting.

Nature Medicine is providing this early version of the typeset paper as a service to our authors and readers. The text and figures will undergo copyediting and a proof review before the paper is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers apply.

41 **Low neutralization of SARS-CoV-2 Omicron BA.2.75.2, BQ.1.1, and XBB.1 by parental**
42 **mRNA vaccine or a BA.5-bivalent booster**

43 Chaitanya Kurhade^{1,#}, Jing Zou^{1,#}, Hongjie Xia¹, Mingru Liu², Hope C. Chang², Ping Ren^{2,*},
44 Xuping Xie^{1,*}, Pei-Yong Shi^{1,3,4,5,6,*}

45 ¹Department of Biochemistry and Molecular Biology, University of Texas Medical Branch,
46 Galveston, TX, USA

47 ²Department of Pathology, University of Texas Medical Branch, Galveston Texas, USA

48 ³Sealy Institute for Drug Discovery, University of Texas Medical Branch, Galveston, TX, USA

49 ⁴Institute for Human Infection and Immunity, University of Texas Medical Branch, Galveston,
50 TX, USA

51 ⁵Sealy Institute for Vaccine Sciences, University of Texas Medical Branch, Galveston, TX, USA

52 ⁶Sealy Center for Structural Biology & Molecular Biophysics, University of Texas Medical
53 Branch, Galveston, TX, USA

54
55 #C.K. and J.Z. made equal contributions to the study.

56 *Corresponding authors: P.R. (peren@utmb.edu), X.X. (xuxie@utmb.edu), or P.-Y.S.
57 (peshi@utmb.edu)

58
59 **Abstract**

60 The newly emerged SARS-CoV-2 Omicron sublineages, including the BA.2-derived
61 BA.2.75.2 and the BA.5-derived BQ.1.1 and XBB.1, have accumulated additional spike mutations
62 that may affect vaccine effectiveness. Here we report neutralizing activities of three human serum
63 panels collected from individuals 23-94 days after dose 4 of a parental mRNA vaccine, 14-32

64 days after a BA.5-bivalent-booster from individuals with 2-4 previous doses of parental mRNA
65 vaccine, or 15-32 days after a BA.5-bivalent-booster from individuals with previous SARS-CoV-2
66 infection and 2-4 doses of parental mRNA vaccine. The results showed that a BA.5-bivalent-
67 booster elicited a high neutralizing titer against BA.4/5 measured at 14- to 32-day post-boost;
68 however, the BA.5-bivalent-booster did not produce robust neutralization against the newly
69 emerged BA.2.75.2, BQ.1.1, or XBB.1. Previous infection significantly enhanced the magnitude
70 and breadth of BA.5-bivalent-booster-elicited neutralization. Our data support a vaccine update
71 strategy that future boosters should match newly emerged circulating SARS-CoV-2 variants.

72

73 **Main text**

74 The continuous emergence of new variants of severe acute respiratory syndrome
75 coronavirus 2 (SARS-CoV-2) has caused successive global waves of infection. Since its first
76 report in November 2021 in South Africa, Omicron has become the dominating variant due to its
77 high transmissibility and immune evasion^{1,2}, with many Omicron sublineages emerging over time.
78 The initial Omicron BA.1 was displaced by BA.2, which has further evolved to sublineages
79 BA.2.12.1, BA.2.75, BA.2.75.2, BA.4, and BA.5, among which BA.5 is currently dominant in many
80 countries. BA.4 and BA.5 have an identical spike sequence (defined as BA.4/5 hereafter) and
81 their offspring BA.4.6, BF.7, and BQ.1.1 are expanding in prevalence. As of November 19, 2022,
82 the BA.2-derived sublineage BA.2.75.2 accounted for 0.8% of the total SARS-CoV-2 infection in
83 the United States; whereas the BA.4/5-derived sublineages BA.4.6, BF.7, BQ.1, and BQ.1.1
84 accounted for 4.4%, 7.8%, 25.5%, and 24.2% of total cases, respectively. In addition, another
85 BA.5-derived sublineage XBB, first identified in India in August 2022, is rapidly spreading in
86 Europe and has been detected in the United States. XBB was predominant in Singapore,
87 accounting for 54% of SARS-CoV-2 infections during the week of October 3-9, 2022 (Ministry of
88 Health, Singapore- <https://www.moh.gov.sg/>).

89 SARS-CoV-2 spike mutations often contribute to immune evasion and/or transmission
90 efficiency³⁻⁹. Previous studies showed that 3 or 4 doses of parental mRNA vaccine did not elicit
91 robust neutralization against BA.4/5, supporting the development of bivalent vaccines that target
92 both the ancestral spike and the BA.4/5 spike protein¹⁰⁻¹². Since the newly emerged Omicron
93 sublineages have accumulated additional spike mutations (**Fig. 1A**), it is important to examine
94 the vaccine-elicited neutralization against these new sublineages. The goal of this study was to
95 compare the neutralizing activities against six newly emerged Omicron sublineages (BA.5, BF.7,
96 BA.4.6, BA.2.75.2, BQ.1.1, and XBB.1) using human sera collected from individuals who received
97 4 doses of parental mRNA vaccine or a BA.5-bivalent-booster after 2-4 doses of parental mRNA
98 vaccine.

99 To facilitate neutralization measurement, we engineered the complete *spike* gene from
100 Omicron sublineage BA.4/5, BF.7, BA.4.6, BA.2.75.2, BQ.1.1, or XBB.1 into the backbone of
101 mNeonGreen (mNG) reporter USA-WA1/2020 SARS-CoV-2 (**Fig. 1A**). Compared with wild-type
102 USA-WA1/2020 (a strain isolated in January 2020), insertion of *mNG* gene at open-reading-
103 frame-7 of the viral genome attenuated the virus *in vivo*¹³. So, the engineered live-attenuated
104 mNG viruses can be used safely in a BSL3 facility with the correct procedures for neutralization
105 and antiviral testing¹⁴. Passage 1 of recombinant BA.4/5-, BF.7-, BA.4.6-, BA.2.75.2-, BQ.1.1-,
106 and XBB.1-spike mNG USA-WA1/2020 viruses were sequenced to ensure no undesired
107 mutations. Only the passage 1 virus stocks were used to determine the 50% fluorescent focus-
108 reduction neutralization titers (FFRNT₅₀) of vaccinated human sera, to ensure no additional spike
109 mutations in the tested recombinant viruses.

110 Three human serum panels with distinct vaccination and/or SARS-CoV-2 infection history
111 were analyzed. The first panel consisted of 25 sera obtained from individuals 23-94 (median 47)
112 days post dose 4 of parental monovalent mRNA-1273 or BNT162b2 vaccine (post-dose-4 sera);
113 these sera were collected from March 16 to June 30, 2022 (Extended Data Table 1). The second

114 panel consisted of 29 sera collected from individuals 14-32 (median 22) days post BA.5-bivalent-
115 booster (BA.5-bivalent-booster sera); these specimens were collected from September 30 to
116 October 22, 2022 (Extended Data Table 2). All sera from the first and second panels tested
117 negative against viral nucleocapsid protein (Extended Data Figure 1), suggesting no previous or
118 recent SARS-CoV-2 infection. The third panel consisted of 23 sera collected from individuals who
119 were previously infected by SARS-CoV-2 (nucleocapsid antibody positive; Extended Data Figure
120 1) and received a BA.5-bivalent-booster 15-32 (median 21) days ago (BA.5-bivalent-booster-
121 infection sera); the viral infection time and genotype could not be determined because most
122 infections were asymptomatic; these samples were collected from October 4 to 22, 2022
123 (Extended Data Table 3). All participants from the second and third panels had also received 2,
124 3, or 4 doses of parental monovalent mRNA vaccine before receiving the BA.5-bivalent-booster.
125 Extended Data Table 1-3 summarize the serum information and neutralization for each serum
126 panel.

127 Post-dose-4 sera neutralized USA-WA1/2020-, BA.4/5-, BF.7-, BA.4.6-, BA.2.75.2-,
128 BQ.1.1-, and XBB.1-spike mNG SARS-CoV-2 with geometric mean titers (GMTs) of 1533, 95, 69,
129 62, 26, 22, and 15, respectively (Figure 1B and Extended Data Table 1). The neutralizing GMTs
130 against BA.4/5-, BF.7-, BA.4.6-, BA.2.75.2-, BQ.1.1-, and XBB.1-spike viruses were 16.1-, 22.2-,
131 24.7-, 59-, 69.7-, and 102-fold lower than the GMT against the USA-WA1/2020-spike virus,
132 respectively (Figure 1B). Compared with the GMT against the current dominant BA.4/5, the
133 neutralizing GMTs against BF.7-, BA.4.6-, BA.2.75.2-, BQ.1.1-, and XBB.1-spike viruses were
134 reduced by 1.4-, 1.5-, 3.7-, 4.3-, and 6.3-fold, respectively. The GMTs against BA.2.75.2 (26) and
135 BQ.1.1 (22) were barely above 20, the detection limit of FFRNT; whereas the GMT against XBB.1
136 (15) was below the FFRNT detection limit. These results indicate that (i) 4 doses of parental
137 mRNA vaccine do not elicit robust neutralization against the newly emerged Omicron sublineages

138 when measured at 23-94 (median 47) days post-dose-4 and (ii) the rank of neutralization evasion
139 is in the order of BA.4/5 < BF.7 ≤ BA.4.6 < BA.2.75.2 ≤ BQ.1.1 < XBB.1.

140 BA.5-bivalent-booster sera, collected at 14-32 (median 22) days post-boost, neutralized
141 USA-WA1/2020-, BA.4/5-, BF.7-, BA.4.6-, BA.2.75.2-, BQ.1.1-, and XBB.1-spike SARS-CoV-2s
142 with GMTs of 3620, 298, 305, 183, 98, 73, and 35, respectively (Figure 1C and Extended Data
143 Table 2). The neutralizing GMTs against BA.4/5-, BF.7-, BA.4.6-, BA.2.75.2-, BQ.1.1-, and XBB.1-
144 spike viruses were 12.1-, 11.9-, 19.8-, 36.9-, 49.6-, and 103-fold lower than the GMT against the
145 USA-WA1/2020, respectively (Figure 1C). The data indicate that although BA.5-bivalent-booster
146 elicits high neutralizing titers against BA.4/5 measured at 14-32 days post-boost, the
147 neutralization against BA.2.75.2 (98), BQ.1.1 (73), and XBB.1 (35) remains low after BA.5-
148 bivalent-booster.

149 BA.5-bivalent-booster-infection sera, collected at 15-32 (median 21) days post-boost,
150 neutralized USA-WA1/2020-, BA.4/5-, BF.7-, BA.4.6-, BA.2.75.2-, BQ.1.1-, and XBB.1-spike
151 SARS-CoV-2s with GMTs of 5776, 1558, 1223, 744, 367, 267, and 103, respectively (Figure 1D
152 and Extended Data Table 3). The neutralizing GMTs against BA.4/5-, BF.7-, BA.4.6-, BA.2.75.2-
153 , BQ.1.1-, and XBB.1-spike viruses were 3.7-, 4.7-, 7.8-, 15.7-, 21.6-, and 56.1-fold lower than the
154 GMT against the USA-WA1/2020-spike SARS-CoV-2, respectively (Figure 1D). Compared with
155 BA.5-bivalent-booster sera without infection history, BA.5-bivalent-booster-infection sera
156 increased the neutralizing GMTs against USA-WA1/2020-, BA.4/5-, BF.7-, BA.4.6-, BA.2.75.2-,
157 BQ.1.1-, and XBB.1-spike viruses by 1.6-, 5.2-, 4.0-, 4.1-, 3.7-, 3.7-, and 2.9-fold, respectively
158 (Compare Figures 1C and 1D). The results suggest that (i) previous infection significantly
159 increases the magnitude and breadth of neutralization for BA.5-bivalent-booster and (ii) among
160 the tested Omicron sublineages, XBB.1 exhibits the highest level of immune evasion.

161 Collectively, our neutralization results support two conclusions. First, the newly emerged
162 Omicron sublineages continue to increase their immune evasion of vaccine- and/or infection-

163 elicited neutralization. Among tested Omicron sublineages, BA.2.75.2, BQ.1.1, and XBB.1 exhibit
164 the greatest evasion against vaccine-elicited neutralization, suggesting the potential of these new
165 sublineages to dethrone BA.5 as the dominant lineage in circulation. Second, individuals with
166 SARS-CoV-2 infection history develop higher and broader neutralization against the current
167 circulating Omicron sublineages after the BA.5-bivalent booster.

168 The study has several limitations. First, we have not examined the antiviral roles of non-
169 neutralizing antibodies and cell-mediated immunity. These two immune components, together with
170 neutralizing antibodies, protect patients from severe disease and death^{15,16}. Unlike neutralizing
171 antibodies, many T cell epitopes after vaccination or natural infection are preserved in Omicron
172 spikes¹⁷. However, robust antibody neutralization is critical to prevent viral infection¹⁸. Second,
173 we have not defined the spike mutations that contribute to the observed immune evasion of the
174 newly emerged Omicron sublineages. Spike mutation F486V was previously shown to drive the
175 immune evasion of BA.4/5¹⁰. The new Omicron sublineages BA.2.75.2, BA.4.6, BF.7, BQ.1.1, and
176 XBB.1 share the spike R346T mutation that was reported to confer higher neutralization
177 evasion¹⁹. Third, the current results do not allow a direct comparison of neutralization between
178 parental mRNA vaccine and BA.5-bivalent-booster because of the differences in individuals'
179 demographics (e.g., age), numbers of vaccine doses, and serum collection time. Fourth, we don't
180 know (i) how neutralizing titers related to protection against infection, severe disease, or death;
181 (ii) when and which variants infected individuals from the BA.5-bivalent-booster-infection cohort;
182 and (iii) the insight related to the differences in vaccine dose for Moderna's Bivalent (Original and
183 Omicron BA. 4/BA. 5) versus Pfizer/BioNTech's BA.4/BA.5-Adapted Bivalent Booster, and (iv) the
184 baseline of the neutralization titers before boost were not determined due to sample unavailability.

185 Our laboratory investigation, along with the recent real-world effectiveness of BA.5-
186 bivalent-booster²⁰, supports a vaccine update strategy that future boosters should match new
187 circulating SARS-CoV-2 variants. Given the advantage of mRNA vaccine platform that can rapidly

188 adapt to new antigen sequences, the key challenge is to determine the future booster sequence
189 before new variants become prevalent in circulation.

190

191 **Acknowledgments**

192 We thank colleagues at the University of Texas Medical Branch (UTMB) for helpful discussions
193 P.-Y.S. was supported by NIH contract HHSN272201600013C, and awards from the Sealy &
194 Smith Foundation, the Kleberg Foundation, the John S. Dunn Foundation, the Amon G. Carter
195 Foundation, the Summerfield Robert Foundation, and Edith and Robert Zinn. We thank the
196 participants from whom the serum specimens were obtained. The funders had no role in study
197 design, data collection and analysis, decision to publish or preparation of the manuscript.

198

199 **Author contributions**

200 Conceptualization, P.R., X.X., P.-Y.S.; Methodology, C.K., J.Z., H.X., M.L., H.C.C., P.R., X.X., P.-
201 Y.S.; Investigation, C.K., J.Z., H.X., M.L., H.C.C., P.R., X.X., P.-Y.S.; Resources, P.R., X.X., P.-
202 Y.S.; Data Curation, C.K., J.Z., P.R., X.X.; Writing-Original Draft, P.R., X.X., P.-Y.S.; Writing-
203 Review & Editing, C.K., J.Z., H.X., M.L., H.C.C., P.R., X.X., P.-Y.S.; Supervision, P.R., X.X., P.-
204 Y.S.; Funding Acquisition, P.R., X.X., P.-Y.S.

205

206 **Competing interests**

207 X.X. and P.-Y.S. have filed a patent on the reverse genetic system. X.X., J.Z., and P.-Y.S.
208 received compensation from Pfizer for COVID-19 vaccine development. Other authors declare
209 no competing interests.

210

211 **Figure Legends**

212

213 Figure 1. Neutralization against Omicron sublineages. (A) Construction of Omicron sublineage-
214 spike mNG SARS-CoV-2. Amino acid mutations, deletions (Δ), and insertions (Ins) are indicated
215 in reference to the USA-WA1/2020 spike. L: leader sequence; ORF: open reading frame; NTD:
216 N-terminal domain of S1; RBD: receptor binding domain; S: spike glycoprotein; S1: N-terminal
217 furin cleavage fragment of S; S2: C-terminal furin cleavage fragment of S; E: envelope protein;
218 M: membrane protein; N: nucleocapsid; UTR: untranslated region. (B) FFRNT₅₀s of human sera
219 after dose 4 parental mRNA vaccine. The p values (two-sided) for group comparison of GMTs are
220 the following. USA-WA1/2020 versus others: <0.0001; BA.4/5-spike versus BF.7-, BA.4.6-,
221 BA.2.75.2-, BQ.1.1-, XBB.1-spike: 0.029, 0.001, <0.0001, <0.0001, <0.0001; BF.7-spike versus
222 BA.4.6-, BA.2.75.2-, BQ.1.1-, XBB.1-spike: 0.103, <0.0001, <0.0001, <0.0001; BA.4.6-spike
223 versus BA.2.75.2-, BQ.1.1-, XBB.1-spike: 0.0001, <0.0001, and <0.0001; BA.2.75.2-spike versus
224 BQ.1.1-, XBB.1-spike: 0.24, <0.0001; BQ.1.1-spike versus XBB.1-spike: 0.0028. The FFRNT₅₀
225 values against BA.4/5-spike were reported previously¹¹. (C) FFRNT₅₀ of 29 sera collected after
226 BA. 5-bivalent booster from individuals without infection history. The p values (two-sided) for
227 group comparison of GMTs are the following. USA-WA1/2020 versus others: <0.0001; BA.4/5-
228 spike versus BF.7-, BA.4.6-, BA.2.75.2-, BQ.1.1-, XBB.1-spike: 0.844, <0.0001, <0.0001,
229 <0.0001, <0.0001; BF.7-spike versus BA.4.6-, BA.2.75.2-, BQ.1.1-, XBB.1-spike: all <0.0001;
230 BA.4.6-spike versus BA.2.75.2-, BQ.1.1-, XBB.1-spike: all <0.0001; BA.2.75.2-spike versus
231 BQ.1.1-, XBB.1-spike: 0.69, <0.0001; BQ.1.1-spike versus XBB.1-spike: <0.0001. (D) FFRNT₅₀
232 of 23 sera collected after BA.5-bivalent-booster from individuals with infection history. The p
233 values (two-sided) for group comparison of GMTs are the following. USA-WA1/2020 versus
234 others: <0.0001; BA.4/5-spike versus BF.7-, BA.4.6-, BA.2.75.2-, BQ.1.1-, XBB.1-spike: 0.0049,
235 <0.0001, <0.0001, <0.0001, <0.0001; BF.7-spike versus BA.4.6-, BA.2.75.2-, BQ.1.1-, XBB.1-
236 spike: all <0.0001; BA.4.6-spike versus BA.2.75.2-, BQ.1.1-, XBB.1-spike: 0.0005, <0.0001,
237 <0.0001; BA.2.75.2-spike versus BQ.1.1-, XBB.1-spike: 0.114, <0.0001; BQ.1.1-spike versus
238 XBB.1-spike: <0.0001. For figure panels b-d, bar heights and the numbers above indicate GMTs.
239 Error bars indicate 95% CI. The fold of GMT reduction against each Omicron sublineage,
240 compared with the GMT against USA-WA1/2020, is shown in italic font. The dotted line indicates
241 the limit of detection of FFRNT₅₀. Statistic analyses were performed using the Wilcoxon matched-
242 pairs signed-rank test for group comparison of GMTs.

243

244 **References**

- 245 1 Frederik Plesner Lyngse *et al.* Transmission of SARS-CoV-2 Omicron VOC subvariants BA.1 and
246 BA.2: Evidence from Danish Households. *BioRxiv*, doi:
247 <https://doi.org/10.1101/2022.1101.1128.22270044> (2022).
- 248 2 Xia, H. *et al.* Neutralization and durability of 2 or 3 doses of the BNT162b2 vaccine against
249 Omicron SARS-CoV-2. *Cell Host Microbe* **30**, 485-488.e483 (2022).
250 <https://doi.org/10.1016/j.chom.2022.02.015>
- 251 3 Zou, J. *et al.* Neutralization against Omicron SARS-CoV-2 from previous non-Omicron infection.
252 *Nat Commun* **13**, 852 (2022). <https://doi.org/10.1038/s41467-022-28544-w>
- 253 4 Liu, Y. *et al.* BNT162b2-Elicited Neutralization against New SARS-CoV-2 Spike Variants. *N Engl J*
254 *Med* **385**, 472-474 (2021). <https://doi.org/10.1056/NEJMc2106083>
- 255 5 Liu, Y. *et al.* Neutralizing Activity of BNT162b2-Elicited Serum. *N Engl J Med* **384**, 1466-1468
256 (2021). <https://doi.org/10.1056/NEJMc2102017>
- 257 6 Plante, J. A. *et al.* Spike mutation D614G alters SARS-CoV-2 fitness. *Nature* **592**, 116-121 (2021).
258 <https://doi.org/10.1038/s41586-020-2895-3>
- 259 7 Liu, Y. *et al.* Delta spike P681R mutation enhances SARS-CoV-2 fitness over Alpha variant. *Cell*
260 *Rep* **39**, 110829 (2022). <https://doi.org/10.1016/j.celrep.2022.110829>
- 261 8 Liu, J. *et al.* BNT162b2-elicited neutralization of B.1.617 and other SARS-CoV-2 variants. *Nature*
262 **596**, :273-275 (2021). <https://doi.org/10.1038/s41586-021-03693-y>
- 263 9 Liu, Y. *et al.* The N501Y spike substitution enhances SARS-CoV-2 transmission. *Nature* **602**, 294-
264 299 (2021). <https://doi.org/10.1101/2021.03.08.434499>
- 265 10 Kurhade, C. *et al.* Neutralization of Omicron sublineages and Deltacron SARS-CoV-2 by 3 doses of
266 BNT162b2 vaccine or BA.1 infection. *Emerg Microbes Infect*, 1-18 (2022).
267 <https://doi.org/10.1080/22221751.2022.2099305>
- 268 11 Xie, X. *et al.* Neutralization of SARS-CoV-2 Omicron sublineages by 4 doses of the original mRNA
269 vaccine. *Cell Rep*, 111729 (2022). <https://doi.org/10.1016/j.celrep.2022.111729>
- 270 12 Kurhade, C. *et al.* Neutralization of Omicron BA.1, BA.2, and BA.3 SARS-CoV-2 by 3 doses of
271 BNT162b2 vaccine. *Nat Commun* **13**, 3602 (2022). <https://doi.org/10.1038/s41467-022-30681-1>
- 272 13 Johnson, B. A. *et al.* Nucleocapsid mutations in SARS-CoV-2 augment replication and
273 pathogenesis. *PLoS Pathog* **18**, e1010627 (2022). <https://doi.org/10.1371/journal.ppat.1010627>
- 274 14 Muruato, A. E. *et al.* A high-throughput neutralizing antibody assay for COVID-19 diagnosis and
275 vaccine evaluation. *Nat Commun* **11**, 4059 (2020). <https://doi.org/10.1038/s41467-020-17892-0>
- 276 15 Bartsch, Y. C. *et al.* Omicron variant Spike-specific antibody binding and Fc activity are preserved
277 in recipients of mRNA or inactivated COVID-19 vaccines. *Sci Transl Med* **14**, eabn9243 (2022).
278 <https://doi.org/10.1126/scitranslmed.abn9243>
- 279 16 Grifoni, A. *et al.* SARS-CoV-2 human T cell epitopes: Adaptive immune response against COVID-
280 19. *Cell Host Microbe* **29**, 1076-1092 (2021). <https://doi.org/10.1016/j.chom.2021.05.010>
- 281 17 Redd, A. D. *et al.* Minimal Crossover between Mutations Associated with Omicron Variant of
282 SARS-CoV-2 and CD8(+) T-Cell Epitopes Identified in COVID-19 Convalescent Individuals. *mBio*,
283 e0361721 (2022). <https://doi.org/10.1128/mbio.03617-21>
- 284 18 Barouch, D. H. Covid-19 Vaccines - Immunity, Variants, Boosters. *N Engl J Med* (2022).
285 <https://doi.org/10.1056/NEJMra2206573>
- 286 19 Jian, F. *et al.* Further humoral immunity evasion of emerging SARS-CoV-2 BA.4 and BA.5
287 subvariants. *Lancet Infect Dis* **22**, 1535-1537 (2022). [https://doi.org/10.1016/S1473-3099\(22\)00642-9](https://doi.org/10.1016/S1473-3099(22)00642-9)
288

289 20 Link-Gelles, R. *et al.* Effectiveness of Bivalent mRNA Vaccines in Preventing Symptomatic SARS-
290 CoV-2 Infection — Increasing Community Access to Testing Program, United States, September–
291 November 2022. *MMWR Morb Mortal Wkly Rep* ePub: **22 November 2022**. (2022).
292 [https://doi.org: http://dx.doi.org/10.15585/mmwr.mm7148e1](https://doi.org/http://dx.doi.org/10.15585/mmwr.mm7148e1)

293

294

ACCELERATED ARTICLE PREVIEW

295

296 **Methods**

297 Ethical statement

298 All virus work was performed in a biosafety level 3 (BSL-3) laboratory with redundant fans in the
299 biosafety cabinets at The University of Texas Medical Branch at Galveston. All personnel wore
300 powered air-purifying respirators (Breathe Easy, 3M) with Tyvek suits, aprons, booties, and
301 double gloves. The research protocol regarding the use of human serum specimens was reviewed
302 and approved by the University of Texas Medical Branch (UTMB) Institutional Review Board (IRB
303 number 20-0070). No informed consent was required since these deidentified sera were leftover
304 specimens from the routine standard of care and diagnostics before being discarded. No
305 diagnosis or treatment was involved either. The use of human serum specimens in this study was
306 reviewed and approved by the University of Texas Medical Branch (UTMB) Institutional Review
307 Board (IRB number 20-0070).

308 Cells

309 Vero E6 (ATCC® CRL-1586) purchased from the American Type Culture Collection (ATCC,
310 Bethesda, MD) and Vero E6 cells expressing TMPRSS2 (JCRB1819) purchased from SEKISUI
311 XenoTech, LLC were maintained in a high-glucose Dulbecco's modified Eagle's medium (DMEM)
312 containing 10% fetal bovine serum (FBS; HyClone Laboratories, South Logan, UT) and 1%
313 penicillin/streptomycin at 37°C with 5% CO₂. Culture media and antibiotics were purchased from
314 Thermo Fisher Scientific (Waltham, MA). Both cell lines were tested negative for *Mycoplasma*.

315 Human Serum

316 Three panels of human sera collected at UTMB were used in the study. Samples were collected
317 based on availability. Varied ages with both genders are included. The population contains varied
318 races or ethnicity, including white, Hispanic, black, and Asian. Subjects have received at least

319 two doses of the COVID-19 vaccine with or without evidence of SARS-CoV-2 infection. The first
320 panel consisted of 25 sera collected from individuals 23-94 (median 47) days after receiving dose
321 4 of parental vaccine mRNA-1273 or BNT162b2. This panel had been tested negative for SARS-
322 CoV-2 nucleocapsid protein expression using Bio-Plex Pro Human IgG SARS-CoV-2
323 N/RBD/S1/S2 4-Plex Panel (Bio-rad). The second panel consisted of 29 sera collected from
324 individuals 14-32 (median 22) days after BA.5-bivalent-booster from Pfizer (BA.4/BA.5-Adapted
325 Bivalent Booster) or Moderna (Bivalent Booster). All sera from this panel were tested negative for
326 antibodies against SARS-CoV-2 nucleocapsid protein. The third panel consisted of 23 sera from
327 individuals who were previously infected by SARS-CoV-2 (as determined by SARS-CoV-2
328 nucleocapsid ELISA), vaccinated with 2-4 doses of parental mRNA vaccine, and received a BA.5-
329 bivalent-booster 15-32 (median 21) days before serum collection. The genotypes of the infecting
330 SARS-CoV-2 variants could not be determined for the third serum panel. Patient information was
331 completely deidentified from all specimens. No informed consent was required because these
332 deidentified sera were leftover specimens from standard care and diagnostics before being
333 discarded. The use of human sera for this study was reviewed and approved by the UTMB IRB
334 (number 20-0070). The de-identified human sera were heat-inactivated at 56°C for 30 min before
335 the neutralization test. The serum information is presented in Extended Data Table 1-3.

336 Generation of recombinant Omicron sublineages-mNG SARS CoV-2

337 Recombinant Omicron sublineage BA.4/5-, BF.7-, BA.4.6-, BA.2.75.2-, BQ.1.1-, and XBB.1-spike
338 mNG SARS-CoV-2s was constructed by engineering the complete *spike* gene from the indicated
339 variants into an infectious cDNA clone of mNG USA-WA1/2020 as reported previously^{21,22}. Spike
340 sequences were based on BA.4/5 (BA.4: GISAID EPI_ISL_11 542270; BA.5: GISAID
341 EPI_ISL_11542604; BA.4 and BA.5 have the identical spike sequence), BA.4.6 (GISAID
342 EPI_ISL_15380489), BA.2.75.2 (GISAID EPI_ISL_14458978), BF.7 (GISAID
343 EPI_ISL_14425795), BQ.1.1 (GISAID EPI_ISL_15542649) and XBB.1 (GISAID

344 EPI_ISL_15232105). The full-length infectious cDNA clone of SARS-CoV-2 was assembled by *in*
345 *vitro* ligation followed by *in vitro* transcription to synthesize the viral genomic RNA. The full-length
346 RNA transcripts were electroporated in Vero E6-TMPRSS2 cells to recover the viruses. Viruses
347 were rescued post 2-3 days after electroporation and served as P0 stock. P0 stock was further
348 passaged once on Vero E6 cells to produce P1 stock. The reason for using Vero E6 cells (rather
349 than using Vero E6-TMPRSS2) to prepare the P1 virus is that the infectivity of the P1 virus can
350 be affected by the cell types; since our established FFRNT assay uses Vero E6 cells, we chose
351 to prepare the P1 viruses using the same Vero E6 cells. The *spike* gene was sequenced from all
352 P1 stock viruses to ensure no undesired mutation. The infectious titer of the P1 virus was
353 quantified by fluorescent focus assay on Vero E6 cells. The P1 virus was used for the
354 neutralization test. The protocols for the mutagenesis of mNG SARS-CoV-2 and virus production
355 were reported previously²³. All virus preparation and neutralization assays were carried out at the
356 biosafety level 3 (BSL-3) facility at the University of Texas Medical Branch at Galveston.

357 Fluorescent focus reduction neutralization test (FFRNT)

358 Neutralization titers of human sera were measured by FFRNT using the USA-WA1/2020-, BA.4/5-
359 , BF.7-, BA.4.6-, BA.2.75.2-, BQ.1.1- and XBB.1-spike mNG SARS-CoV-2s at BSL-3. The details
360 of the FFRNT protocol were reported previously²³. Briefly, 2.5×10^4 Vero E6 cells per well were
361 seeded in 96-well plates (Greiner Bio-one™). The cells were incubated overnight. On the next
362 day, each serum was 2-fold serially diluted in the culture medium with the first dilution of 1:20
363 (final dilution range of 1:20 to 1:20,480). The diluted serum was incubated with 100-150 FFUs of
364 mNG SARS-CoV-2 at 37 °C for 1 h, after which the serum virus mixtures were loaded onto the
365 pre-seeded Vero E6 cell monolayer in 96-well plates. After 1 h infection, the inoculum was
366 removed and 100 µl of overlay medium (supplemented with 0.8% methylcellulose) was added to
367 each well. After incubating the plates at 37 °C for 16 h, raw images of mNG foci were acquired
368 using Cytation™ 7 (BioTek) armed with 2.5x FL Zeiss objective with a wide-field of view and

369 processed using the Gen 5 software settings (GFP [469,525] threshold 4000, object selection size
370 50-1000 μm). The foci in each well were counted using the Gen5 software and normalized to the
371 non-serum-treated controls to calculate the relative infectivities. The FFRNT₅₀ value was defined
372 as the minimal serum dilution that suppressed >50% of fluorescent foci. The neutralization titer of
373 each serum was determined in duplicate assays, and the geometric mean was taken. Tables S1-
374 3 summarize the FFRNT₅₀ results. Data were initially plotted in GraphPad Prism 9 software and
375 assembled in Adobe Illustrator. FFRNT₅₀ of <20 was treated as 10 for plot purposes and
376 statistical analysis. The above FFRNT₅₀ protocol has been reliably used to support the clinical
377 development of COVID-19 vaccines²⁴. Thus, we applied the same FFRNT protocol to the current
378 study.

379 Statistics & Reproducibility

380 No statistical method was used to predetermine the sample size. The samples were collected
381 based on availability. No data were excluded from the analyses. The experiments were not
382 randomized. Patient information was blinded in the study. The investigators were blinded to
383 sample identity during data collection and/or analysis. The experiments were performed in
384 duplication. All attempts at replication were successful.





385 Continuous variables were summarized as the geometric mean with 95% confidence intervals or
386 median. Sera with undetectable (<20) antibody titers were assigned an antibody titer of 10, for
387 purposes of GMT calculations or statistical comparisons. Comparison between neutralization
388 titers was performed using a Wilcoxon matched-pairs signed-rank test using GraphPad Prism
389 9.0. Absolute *P* values were provided. *P* < 0.05 was considered statistically significant. Images
390 were assembled using Adobe Illustrator.

391 **Data availability**

392 The raw data that support the findings of this study are shown in the Source data files. The
393 sequence of SARS-CoV-2 variants can be accessed through GISAID (<https://gisaid.org>) with the
394 following codes: BA.4/5 (BA.4: GISAID EPI_ISL_11 542270; BA.5: GISAID EPI_ISL_11542604;
395 BA.4 and BA.5 have the identical spike sequence), BA.4.6 (GISAID EPI_ISL_15380489),
396 BA.2.75.2 (GISAID EPI_ISL_14458978), BF.7 (GISAID EPI_ISL_14425795), BQ.1.1 (GISAID
397 EPI_ISL_15542649) and XBB.1 (GISAID EPI_ISL_15232105). The sequence of SARS-CoV-2
398 mNG can be found in the supplementary information of our previous study²⁵.


399 Methods-only reference

- 400 21 Xie, X. *et al.* An Infectious cDNA Clone of SARS-CoV-2. *Cell Host Microbe* **27**, 841-848 e843
401 (2020). <https://doi.org/10.1016/j.chom.2020.04.004>
402 22 Xie, X. *et al.* Engineering SARS-CoV-2 using a reverse genetic system. *Nat Protoc* **16**, 1761-1784
403 (2021). <https://doi.org/10.1038/s41596-021-00491-8>
404 23 Xie, X. *et al.* Neutralization of SARS-CoV-2 Omicron sublineages by 4 doses of the original mRNA
405 vaccine. *Cell Reports* **41**, 111729 (2022).
406 <https://doi.org/10.1016/j.celrep.2022.111729>
407 24 Mulligan, M. J. *et al.* Phase I/II study of COVID-19 RNA vaccine BNT162b1 in adults. *Nature* **586**,
408 589-593 (2020). <https://doi.org/10.1038/s41586-020-2639-4>
409 25 Muruato, A. E. *et al.* A high-throughput neutralizing antibody assay for COVID-19 diagnosis and
410 vaccine evaluation. *Nat Commun* **11**, 4059 (2020). <https://doi.org/10.1038/s41467-020-17892-0>



ACCELERATING ARTICLES PREVIEW

Nucleocapsid IgG



| Serum ID | Age (year) | Gender (F/M) | Race or Ethnicity | Serum collection time (days post-dose 4 vaccination) | Serum collection date | mRNA Vaccine type | *FFRNT ₅₀ | | | | | | |
|---------------------|------------|--------------|-------------------|--|-----------------------|--|----------------------|---------------------------|------------|--------------|-----------------|--------------|-------------|
| | | | | | | | USA-WA1/2020 | [‡] BA.4/5-spike | BF.7-spike | BA.4.6-spike | BA.2.75.2-spike | BQ.1.1-spike | XBB.1-spike |
| 1 | 62 | F | White | 37 | 3/16/2022 | BTN162b | 320 | [^] 10 | 10 | 10 | 10 | 10 | 10 |
| 2 | 80 | M | Hispanic | 30 | 4/7/2022 | BTN162b | 10240 | 320 | 80 | 80 | 40 | 20 | 20 |
| 3 | 84 | F | White | 55 | 5/11/2022 | BTN162b | 1280 | 80 | 14 | 10 | 10 | 10 | 10 |
| 4 | 92 | F | White | 27 | 5/12/2022 | BTN162b | 2560 | 113 | 40 | 40 | 28 | 10 | 10 |
| 5 | 78 | M | White | 56 | 5/17/2022 | BTN162b | 320 | 10 | 10 | 14 | 10 | 10 | 10 |
| 6 | 71 | F | White | 27 | 5/17/2022 | mRNA-1273 (dose 1-3), BTN162b (dose 4) | 2560 | 320 | 320 | 320 | 80 | 80 | 40 |
| 7 | 83 | F | White | 23 | 5/18/2022 | BTN162b | 1280 | 160 | 113 | 80 | 20 | 20 | 10 |
| 8 | 87 | M | White | 24 | 5/19/2022 | BTN162b | 1280 | 113 | 226 | 160 | 10 | 57 | 10 |
| 9 | 80 | M | White | 52 | 5/23/2022 | BTN162b | 5120 | 640 | 320 | 226 | 80 | 80 | 40 |
| 10 | 84 | M | White | 26 | 5/24/2022 | BTN162b | 640 | 20 | 10 | 10 | 10 | 10 | 10 |
| 11 | 75 | M | Black | 47 | 5/25/2022 | BTN162b | 1810 | 57 | 40 | 40 | 226 | 20 | 40 |
| 12 | 90 | M | Black | 34 | 5/25/2022 | BTN162b | 2560 | 160 | 80 | 80 | 40 | 40 | 20 |
| 13 | 59 | F | Hispanic | 27 | 5/25/2022 | BTN162b (dose 1-3), mRNA-1273 (dose 4) | 1810 | 160 | 113 | 160 | 10 | 28 | 10 |
| 14 | 72 | F | White | 52 | 6/2/2022 | BTN162b | 1810 | 40 | 40 | 40 | 20 | 14 | 10 |
| 15 | 73 | M | White | 94 | 6/3/2022 | mRNA-1273 | 640 | 40 | 40 | 28 | 10 | 10 | 10 |
| 16 | 67 | F | Black | 50 | 6/7/2022 | BTN162b | 1280 | 80 | 80 | 80 | 20 | 40 | 10 |
| 17 | 75 | F | White | 78 | 6/8/2022 | BTN162b | 2560 | 453 | 320 | 320 | 80 | 80 | 40 |
| 18 | 86 | M | White | 49 | 6/9/2022 | BTN162b | 1810 | 320 | 226 | 320 | 40 | 20 | 20 |
| 19 | 66 | F | White | 48 | 6/9/2022 | BTN162b | 453 | 20 | 20 | 14 | 10 | 10 | 10 |
| 20 | 80 | M | Black | 44 | 6/10/2022 | BTN162b | 1280 | 160 | 160 | 160 | 40 | 28 | 20 |
| 21 | 78 | M | White | 73 | 6/13/2022 | BTN162b | 905 | 40 | 40 | 28 | 20 | 10 | 10 |
| 22 | 86 | M | White | 51 | 6/16/2022 | BTN162b | 1280 | 20 | 20 | 20 | 10 | 10 | 10 |
| 23 | 84 | F | White | 35 | 6/20/2022 | BTN162b | 20480 | 1280 | 640 | 453 | 226 | 160 | 80 |
| 24 | 94 | F | White | 47 | 6/26/2022 | BTN162b | 1280 | 80 | 40 | 40 | 20 | 20 | 10 |
| 25 | 87 | F | Hispanic | 43 | 6/30/2022 | BTN162b | 905 | 160 | 320 | 160 | 28 | 14 | 10 |
| Median | 80 | - | - | 47 | - | - | - | - | - | - | - | - | - |
| [#] GMT | - | - | - | - | - | - | 1533 | 95 | 69 | 62 | 26 | 22 | 15 |
| [†] 95% CI | - | - | - | - | - | - | 1036-2268 | 56-160 | 41-115 | 38-103 | 17-38 | 16-31 | 12-20 |

*Individual FFRNT₅₀ value is the geometric mean of duplicate FFRNT results.

[^]FFRNT₅₀ of <20 was treated as 10 for plot purposes and statistical analysis.

[#]Geometric mean neutralizing titers (GMT).

[†]95% confidence interval (95% CI) for the GMT.

[§]This data set was reported previously⁵.

| Serum ID | Age (year) | Gender (F/M) | Race or Ethnicity | Serum collection time (days post-BA.5-bivalent booster) | Serum collection date | Last dose of mRNA vaccine before BA.5-bivalent booster | *FFRNT ₅₀ | | | | | | |
|----------|------------|--------------|-------------------|---|-----------------------|--|----------------------|--------------|------------|--------------|-----------------|--------------|-------------|
| | | | | | | | USA-WA1/2020 | BA.4/5-spike | BF.7-spike | BA.4.6-spike | BA.2.75.2-spike | BQ.1.1-spike | XBB.1-spike |
| 1 | 34 | F | Black | 15 | 9/30/2022 | Dose 3 | 5120 | 640 | 1280 | 640 | 640 | 226 | 160 |
| 2 | 78 | F | White | 17 | 10/3/2022 | Dose 4 | 5120 | 640 | 640 | 320 | 320 | 160 | 80 |
| 3 | 86 | M | White | 21 | 10/4/2022 | Dose 4 | 1280 | 28 | 10 | 10 | 20 | 10 | 10 |
| 4 | 31 | M | Asian | 20 | 10/6/2022 | Dose 2 | 5120 | 1280 | 1280 | 905 | 160 | 160 | 80 |
| 5 | 61 | F | White | 15 | 10/11/2022 | Dose 4 | 5120 | 320 | 453 | 320 | 160 | 80 | 57 |
| 6 | 58 | F | Black | 14 | 10/11/2022 | Dose 3 | 7241 | 1810 | 1810 | 1280 | 320 | 640 | 113 |
| 7 | 69 | M | White | 22 | 10/11/2022 | Dose 4 | 2560 | 80 | 80 | 57 | 57 | 20 | 14 |
| 8 | 67 | F | Asian | 21 | 10/12/2022 | Dose 4 | 20480 | 2560 | 1280 | 640 | 160 | 160 | 40 |
| 9 | 77 | M | Asian | 27 | 10/13/2022 | Dose 4 | 2560 | 226 | 320 | 160 | 80 | 40 | 40 |
| 10 | 39 | F | White | 15 | 10/13/2022 | Dose 3 | 5120 | 320 | 320 | 226 | 80 | 160 | 20 |
| 11 | 73 | M | White | 24 | 10/13/2022 | Dose 4 | 1810 | 160 | 160 | 80 | 80 | 40 | 28 |
| 12 | 83 | F | White | 17 | 10/13/2022 | Dose 4 | 1280 | 40 | 80 | 40 | 40 | 10 | 10 |
| 13 | 79 | F | White | 26 | 10/16/2022 | Dose 3 | 3620 | 320 | 640 | 320 | 226 | 57 | 57 |
| 14 | 35 | F | Asian | 29 | 10/21/2022 | Dose 3 | 5120 | 160 | 320 | 226 | 80 | 80 | 40 |
| 15 | 73 | M | White | 26 | 10/17/2022 | Dose 3 | 7241 | 320 | 320 | 160 | 80 | 80 | 28 |
| 16 | 76 | M | White | 32 | 10/17/2022 | Dose 4 | 7241 | 1280 | 905 | 640 | 226 | 160 | 160 |
| 17 | 71 | M | White | 28 | 10/18/2022 | Dose 4 | 1280 | 40 | 57 | 40 | 40 | 20 | 10 |
| 18 | 22 | M | Hispanic | 19 | 10/18/2022 | Dose 3 | 10240 | 2560 | 1280 | 1280 | 320 | 453 | 80 |
| 19 | 61 | F | White | 30 | 10/19/2022 | Dose 4 | 640 | 80 | 80 | 57 | 20 | 28 | 14 |
| 20 | 56 | M | White | 21 | 10/19/2022 | Dose 3 | 10240 | 640 | 640 | 453 | 160 | 160 | 80 |
| 21 | 66 | F | White | 26 | 10/19/2022 | Dose 4 | 3620 | 320 | 320 | 160 | 80 | 80 | 40 |
| 22 | 76 | F | White | 30 | 10/20/2022 | Dose 4 | 3620 | 226 | 226 | 160 | 10 | 80 | 10 |
| 23 | 61 | F | Asian | 31 | 10/20/2022 | Dose 3 | 10240 | 5120 | 5120 | 2560 | 640 | 453 | 226 |
| 24 | 77 | F | Black | 31 | 10/20/2022 | Dose 4 | 5120 | 80 | 80 | 40 | 40 | 28 | 10 |
| 25 | 59 | F | unknown | 28 | 10/21/2022 | Dose 4 | 2560 | 320 | 320 | 160 | 160 | 80 | 28 |
| 26 | 71 | M | Hispanic | 22 | 10/21/2022 | Dose 4 | 1280 | 160 | 160 | 80 | 80 | 80 | 14 |
| 27 | 70 | F | White | 22 | 10/21/2022 | Dose 3 | 1280 | 113 | 57 | 28 | 20 | 20 | 10 |
| 28 | 79 | F | White | 25 | 10/22/2022 | Dose 4 | 2560 | 226 | 160 | 80 | 57 | 10 | 20 |
| 29 | 79 | M | White | 18 | 10/22/2022 | Dose 4 | 2560 | 160 | 320 | 160 | 226 | 160 | 80 |
| Median | 70 | - | - | 22 | - | - | - | - | - | - | - | - | - |
| *GMT | - | - | - | - | - | - | 3620 | 298 | 305 | 183 | 98 | 73 | 35 |
| †95% CI | - | - | - | - | - | - | 2668-4912 | 181-490 | 185-503 | 111-299 | 66-146 | 47-112 | 24-50 |

*Individual FFRNT₅₀ value is the geometric mean of duplicate FFRNT results.

#Geometric mean neutralizing titers (GMT).

†95% confidence interval (95% CI) for the GMT.

| Serum ID | Age (year) | Gender (F/M) | Race or Ethnicity | Serum collection time (days post-BA.5-bivalent booster) | Serum collection date | Last dose of mRNA vaccine before BA.5-bivalent booster | *FFRNT ₅₀ | | | | | | |
|----------|------------|--------------|-------------------|---|-----------------------|--|----------------------|--------------|------------|--------------|-----------------|--------------|-------------|
| | | | | | | | USA-WA1/2020 | BA.4/5-spike | BF.7-spike | BA.4.6-spike | BA.2.75.2-spike | BQ.1.1-spike | XBB.1-spike |
| 1 | 19 | F | White | 19 | 10/4/2022 | Dose 3 | 10240 | 1280 | 1280 | 640 | 640 | 160 | 113 |
| 2 | 69 | F | White | 15 | 10/6/2022 | Dose 4 | 10240 | 5120 | 5120 | 2560 | 905 | 1280 | 320 |
| 3 | 59 | F | White | 15 | 10/7/2022 | Dose 3 | 20480 | 10240 | 10240 | 7241 | 2560 | 2560 | 453 |
| 4 | 46 | F | White | 18 | 10/11/2022 | Dose 2 | 5120 | 2560 | 2560 | 640 | 320 | 320 | 160 |
| §5 | 80 | F | Black | 22 | 10/11/2022 | Dose 3 | 10240 | 7241 | 5120 | 3620 | 2560 | 640 | 320 |
| 6 | 76 | F | Hispanic | 21 | 10/12/2022 | Dose 3 | 5120 | 1280 | 1280 | 905 | 453 | 160 | 113 |
| 7 | 67 | M | White | 15 | 10/12/2022 | Dose 4 | 5120 | 2560 | 1280 | 1280 | 320 | 320 | 80 |
| 8 | 52 | F | White | 30 | 10/13/2022 | Dose 4 | 5120 | 1280 | 1280 | 905 | 160 | 640 | 160 |
| 9 | 48 | F | Asian | 24 | 10/13/2022 | Dose 3 | 14482 | 7241 | 5120 | 2560 | 1280 | 640 | 320 |
| 10 | 58 | F | White | 28 | 10/13/2022 | Dose 2 | 5120 | 640 | 640 | 320 | 160 | 226 | 40 |
| 11 | 67 | F | Black | 25 | 10/14/2022 | Dose 2 | 5120 | 2560 | 2560 | 1280 | 640 | 640 | 160 |
| 12 | 20 | F | Hispanic | 21 | 10/14/2022 | Dose 2 | 1810 | 320 | 453 | 160 | 80 | 160 | 28 |
| §13 | 75 | M | White | 21 | 10/14/2022 | Dose 3 | 20480 | 2560 | 2560 | 1280 | 1280 | 453 | 320 |
| 14 | 64 | F | White | 19 | 10/4/2022 | Dose 4 | 7241 | 2560 | 2560 | 1280 | 2560 | 453 | 320 |
| 15 | 90 | F | White | 22 | 10/18/2022 | Dose 3 | 10240 | 1280 | 1280 | 1280 | 320 | 640 | 160 |
| 16 | 35 | F | Hispanic | 32 | 10/18/2022 | Dose 2 | 1280 | 320 | 226 | 160 | 80 | 80 | 28 |
| §17 | 39 | M | Asian | 31 | 10/19/2022 | Dose 3 | 10240 | 5120 | 2560 | 2560 | 640 | 320 | 80 |
| §18 | 67 | M | White | 17 | 10/19/2022 | Dose 4 | 10240 | 2560 | 1810 | 1280 | 640 | 640 | 226 |
| §19 | 68 | M | Hispanic | 14 | 10/19/2022 | Dose 3 | 5120 | 2560 | 1280 | 905 | 320 | 160 | 80 |
| 20 | 51 | M | White | 28 | 10/21/2022 | Dose 3 | 7241 | 905 | 320 | 320 | 320 | 160 | 57 |
| 21 | 65 | M | Asian | 29 | 10/20/2022 | Dose 4 | 2560 | 453 | 160 | 80 | 113 | 57 | 40 |
| 22 | 64 | M | White | 16 | 10/21/2022 | Dose 3 | 2560 | 640 | 640 | 320 | 40 | 40 | 20 |
| §23 | 73 | M | Black | 31 | 10/22/2022 | Dose 4 | 640 | 57 | 40 | 28 | 20 | 10 | 10 |
| Median | 64 | - | - | 21 | - | - | - | - | - | - | - | - | - |
| *GMT | - | - | - | - | - | - | 5776 | 1558 | 1223 | 744 | 367 | 267 | 103 |
| †95% CI | - | - | - | - | - | - | 3994-8352 | 922-2631 | 704-2125 | 425-1301 | 209-644 | 158-452 | 66-162 |

*Individual FFRNT₅₀ value is the geometric mean of duplicate FFRNT results.

#Geometric mean neutralizing titers (GMT).

†95% confidence interval (95% CI) for the GMT.

§Reported as SARS-CoV-2 RT-PCR positive. Nucleocapsid testing was not performed.

Reporting Summary

Nature Portfolio wishes to improve the reproducibility of the work that we publish. This form provides structure for consistency and transparency in reporting. For further information on Nature Portfolio policies, see our [Editorial Policies](#) and the [Editorial Policy Checklist](#).

Statistics

For all statistical analyses, confirm that the following items are present in the figure legend, table legend, main text, or Methods section.

n/a Confirmed

- The exact sample size (n) for each experimental group/condition, given as a discrete number and unit of measurement
- A statement on whether measurements were taken from distinct samples or whether the same sample was measured repeatedly
- The statistical test(s) used AND whether they are one- or two-sided
Only common tests should be described solely by name; describe more complex techniques in the Methods section.
- A description of all covariates tested
- A description of any assumptions or corrections, such as tests of normality and adjustment for multiple comparisons
- A full description of the statistical parameters including central tendency (e.g. means) or other basic estimates (e.g. regression coefficient) AND variation (e.g. standard deviation) or associated estimates of uncertainty (e.g. confidence intervals)
- For null hypothesis testing, the test statistic (e.g. F , t , r) with confidence intervals, effect sizes, degrees of freedom and P value noted
Give P values as exact values whenever suitable.
- For Bayesian analysis, information on the choice of priors and Markov chain Monte Carlo settings
- For hierarchical and complex designs, identification of the appropriate level for tests and full reporting of outcomes
- Estimates of effect sizes (e.g. Cohen's d , Pearson's r), indicating how they were calculated

Our web collection on [statistics for biologists](#) contains articles on many of the points above.

Software and code

Policy information about [availability of computer code](#)

Data collection

Data analysis

For manuscripts utilizing custom algorithms or software that are central to the research but not yet described in published literature, software must be made available to editors and reviewers. We strongly encourage code deposition in a community repository (e.g. GitHub). See the Nature Portfolio [guidelines for submitting code & software](#) for further information.

Data

Policy information about [availability of data](#)

All manuscripts must include a [data availability statement](#). This statement should provide the following information, where applicable:

- Accession codes, unique identifiers, or web links for publicly available datasets
- A description of any restrictions on data availability
- For clinical datasets or third party data, please ensure that the statement adheres to our [policy](#)

The raw data that support the findings of this study are shown in the Source data files. The sequence of SARS-CoV-2 variants can be accessed through GISAID (<https://gisaid.org>) with the following codes: BA.4/5 (BA.4: GISAID EPI_ISL_11 542270; BA.5: GISAID EPI_ISL_11542604; BA.4 and BA.5 have the identical spike sequence), BA.4.6 (GISAID EPI_ISL_15380489), BA.2.75.2 (GISAID EPI_ISL_14458978), BF.7 (GISAID EPI_ISL_14425795), BQ.1.1 (GISAID EPI_ISL_15542649) and XBB.1 (GISAID EPI_ISL_15232105). The sequence of SARS-CoV-2 mNG can be found in the supplementary information of our previous study.

Human research participants

Policy information about [studies involving human research participants and Sex and Gender in Research](#).

| | |
|-----------------------------|---|
| Reporting on sex and gender | Human serum samples were collected based on availability. Both gender are included. |
| Population characteristics | Samples were collected based on availability. Varied ages with both genders are included. The population contains varied races or ethnicity, including white, Hispanic, black, and Asian. Subjects have received at least two doses of the COVID-19 vaccine with or without evidence of SARS-CoV-2 infection. |
| Recruitment | No patients were recruited in this study. No informed consent was required, because these deidentified sera were leftover specimens before being discarded. No diagnoses or treatment was involved either. |
| Ethics oversight | The research protocol regarding the use of human serum specimens was reviewed and approved by the University of Texas Medical Branch (UTMB) Institutional Review Board (IRB#: 20-0070). |

Note that full information on the approval of the study protocol must also be provided in the manuscript.

Field-specific reporting

Please select the one below that is the best fit for your research. If you are not sure, read the appropriate sections before making your selection.

Life sciences Behavioural & social sciences Ecological, evolutionary & environmental sciences

For a reference copy of the document with all sections, see [nature.com/documents/nr-reporting-summary-flat.pdf](https://www.nature.com/documents/nr-reporting-summary-flat.pdf)

Life sciences study design

All studies must disclose on these points even when the disclosure is negative.

| | |
|-----------------|---|
| Sample size | No sample size calculation was performed. Samples were collected based on the availability. |
| Data exclusions | No data was excluded in the study. |
| Replication | Each human serum sample was analyzed in duplication. The averaged results from the duplication were reported in this study. All attempts at replication were successful. |
| Randomization | This is no randomization in this study. All samples available were analyzed for the neutralizing activities against WT SARS-CoV-2 and variants in the same experimental settings. |
| Blinding | Patient information was blinded in the study. The investigators were blinded to sample identity during data collection and/or analysis. |

Reporting for specific materials, systems and methods

We require information from authors about some types of materials, experimental systems and methods used in many studies. Here, indicate whether each material, system or method listed is relevant to your study. If you are not sure if a list item applies to your research, read the appropriate section before selecting a response.

Materials & experimental systems

| n/a | Involvement in the study |
|-------------------------------------|---|
| <input checked="" type="checkbox"/> | <input type="checkbox"/> Antibodies |
| <input type="checkbox"/> | <input checked="" type="checkbox"/> Eukaryotic cell lines |
| <input checked="" type="checkbox"/> | <input type="checkbox"/> Palaeontology and archaeology |
| <input checked="" type="checkbox"/> | <input type="checkbox"/> Animals and other organisms |
| <input checked="" type="checkbox"/> | <input type="checkbox"/> Clinical data |
| <input checked="" type="checkbox"/> | <input type="checkbox"/> Dual use research of concern |

Methods

| n/a | Involvement in the study |
|-------------------------------------|---|
| <input checked="" type="checkbox"/> | <input type="checkbox"/> ChIP-seq |
| <input checked="" type="checkbox"/> | <input type="checkbox"/> Flow cytometry |
| <input checked="" type="checkbox"/> | <input type="checkbox"/> MRI-based neuroimaging |

Eukaryotic cell lines

Policy information about [cell lines and Sex and Gender in Research](#)

| | |
|---|---|
| Cell line source(s) | Vero E6 cells (ATCC® CRL-1586) were obtained from ATCC; VeroE6-TMPRSS2 cells were Vero E6 cells expressing TMPRSS2 purchased from SEKISUI XenoTech, LLC. |
| Authentication | VeroE6 Cells have been authenticated by ATCC using morphologies and other groups using STR profiling (reference: Almeida JL, Hill CR, Cole KD. VeroE6-TMPRSS2 cells have been authenticated by the vendor. Authentication of African green monkey cell lines using human short tandem repeat markers. BMC Biotechnol. 2011;11:102. Published 2011 Nov 7. doi:10.1186/1472-6750-11-102). |
| Mycoplasma contamination | All cell lines were tested negative for mycoplasma. |
| Commonly misidentified lines (See ICLAC register) | None. |